

Will Pop Star Troye Sivan Find Love in the HIV Movie “Three Months”?

[VIDEO]

The coming-of-age film centers on HIV test results and also stars Javier Muñoz, Ellen Burstyn and Louis Gossett Jr. Watch the trailer!

February 14, 2022 By Trent Straube

The day before his high school graduation, queer South Florida youth Caleb—who loves his camera, his weed and his grandmother—finds out he’s been exposed to [HIV](#). As he waits 90 days for the definitive results of his [HIV test](#), Caleb finds [love](#) in the most unlikely of places.

That’s the hook for the upcoming movie Three Months (you can watch the trailer above and [on YouTube](#)). The film stars pop crooner Troye Sivan alongside Oscar winners Ellen Burstyn and Louis Gossett Jr. as well as Broadway star Javier Muñoz of Hamilton and In the Heights fame, who in real life is HIV positive.



[View this post on Instagram](#)

A post shared by troye sivan (@troyesivan)

Written and directed by Jared Frieder and produced by MTV Entertainment Studios, *Three Months* will stream exclusively on Paramount+ beginning Wednesday, February 23. It also stars Viveik Kalra (*Blinded by the Light*), Judy Greer (*Halloween*) and Amy Landecker (*Transparent*) and features two new tracks by Sivan.

“Accidental lover, what’d you do with all my common sense?” Sivan questions in one of the soundtrack’s offerings before adding: “I’m tired of living like I’m dying / Now I can find the silver lining.”

From left: Viveik Kalra and Troye Sivan star in “*Three Months*.”MTV Entertainment Studios/Paramount+

In the 2018 film *Boy Erased*, Sivan, who came out as gay in 2013, played a young man at a gay conversion therapy camp. In 2016, Muñoz shared his HIV journey with *POZ* in the cover story “[In the Spotlight](#).”

Not having seen the film—yet!—we can’t say whether the plot follows the latest science about HIV transmission or how it presents life with HIV. But in an article in [Entertainment Weekly](#), Frieder offered promising statements: “At its core, it’s a film about dismantling the shame around HIV and

showing that it's no longer a death sentence in our modern world with access to health care. It's about a kid coming of age and realizing that the very things that make him different also make him indispensable and worthy of love. And perhaps most of all, it's about what it's like to wait."

Troye Sivan and Brianne Tju star in "Three Months."MTV Entertainment Studios/Paramount+

With the advent of rapid HIV tests, you might wonder whether in 2022, a person must wait three months for a test result. The answer is: It depends on the HIV test. As the [POZ Basics on HIV Testing](#) explains:

When it comes to HIV testing, the old cliché “knowledge is power” still holds true. Knowing your HIV status, whether negative or positive, puts you in the best position to protect your health and the health of your sexual partners. While there is no cure for HIV or AIDS, effective medications allow HIV-positive people to live a normal, healthy life. Being aware of your HIV status makes it less likely that you will pass the virus to others, both because you and your partners can take precautions and because effective treatment prevents transmission....

What are the types of HIV tests?

There are several different tests that can be used to determine if you have HIV: antibody tests, antigen/antibody tests and nucleic acid tests. None of these tests are able to detect HIV right away. The time between HIV exposure and when the test can detect the virus is referred to as the window period.

- Antibody tests look for antibodies to HIV in your blood or oral fluid; the ELISA test is most widely used. This is the most common type of HIV screening test. Antibody tests can take 23 to 90 days to detect HIV infection after an exposure because the body takes time to produce antibodies. Most rapid tests and the only Food and Drug Administration–approved HIV self-test are antibody tests. In general, antibody tests that use blood from a vein can detect HIV sooner after infection than tests done with blood from a finger prick or an oral fluid sample.
- Antigen/antibody tests look for both HIV antibodies and antigens, or proteins from the virus itself. An HIV antigen called p24 becomes detectable in the blood before antibodies are produced. Antigen/antibody tests are done in a laboratory. A test using blood from a vein can usually detect HIV infection 18 to 45 days after exposure, while those done using a drop of blood from a finger prick can take longer (18 to 90 days).
- Nucleic acid tests look for HIV genetic material (typically RNA) in the blood. This laboratory test can usually detect HIV infection 10 to 33 days after exposure. The test is expensive and not used for routine screening, but it may be used in some cases to detect early infection. This is the same type of test used to measure viral load and guide treatment for people living with HIV.

Many testing sites now use rapid oral or finger-prick tests. The oral test involves swabbing the upper and lower gums inside the mouth. The sample is then placed into a developer vial, with results available within 20 to 40 minutes. Home HIV self-testing kits are also available.

If an initial rapid test or self-test result is positive, it is important to see a health care provider for a follow-up laboratory test. If the initial HIV test was done through a lab, follow-up testing can usually be done using the same blood sample. HIV tests are generally accurate, but follow-up testing can confirm the diagnosis. If you get an HIV test after a potential exposure and the result is negative, get tested again after the window period end.

Although youth experienced an overall decrease in new [#HIV](#) diagnoses from 2012 to 2019, they still represented over 20% of all new 2019 HIV diagnoses.

Read more this [#NYHAAD: https://t.co/eFdz4K01rapic.twitter.com/bC4mf9p2oG](https://t.co/eFdz4K01rapic.twitter.com/bC4mf9p2oG)
— AIDSvu (@AIDSvu) [April 8, 2021](#)

Visit the [Basics on HIV/AIDS](#) for more details about testing, which also includes the sections [HIV Transmission and Risk](#), [Starting HIV Treatment](#) and [Newly Diagnosed](#).

Did you know youth ages 13 to 24 accounted for over 20% of new HIV cases in 2019? To learn more about HIV in this population, click the hashtag [#Youth](#). You'll find headlines such as:

- [“When Should Youth Get Tested for HIV and Start PrEP?”](#)
- [“Give Young People the Sex Education They Deserve”](#)
- [“National Youth HIV & AIDS Awareness Day”](#)

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.celebhealth.com/article/will-pop-star-troye-sivan-find-love-hiv-movie-three-months-video>